

Observation of e^+e^- Annihilations into the $C = +1$ Hadronic Final States $\rho^0\rho^0$ and $\phi\rho^0$

B. Aubert,¹ R. Barate,¹ M. Bona,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹
V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴
G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶
E. Charles,⁶ M. S. Gill,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶
G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶
W. A. Wenzel,⁶ P. del Amo Sanchez,⁷ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷
S. E. Morgan,⁷ A. T. Watson,⁷ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸
T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ T. Cuhadar-Donszelmann,¹⁰
B. G. Fulsom,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹
M. Saleem,¹¹ D. J. Sherwood,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹²
V. B. Golubev,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu Todyshev,¹²
D. S. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³
P. Lund,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴
S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶
H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷
T. M. Hong,¹⁷ D. Kovalskyi,¹⁷ J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸
J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸
D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ A. Dvoretzskii,¹⁹ F. Fang,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹
T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰
F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹
W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²²
A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² F. Winklmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³
A. Hauke,²³ H. Jasper,²³ A. Petzold,²³ B. Spaan,²³ T. Brandt,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ W. F. Mader,²⁴
R. Nogowski,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵
G. R. Bonneaud,²⁵ P. Grenier,^{25,*} E. Latour,²⁵ Ch. Thiebaux,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶
W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ A. I. Robertson,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷
R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ A. Petrella,²⁷ L. Piemontese,²⁷ E. Prencipe,²⁷
F. Anulli,²⁸ R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸
I. M. Peruzzi,^{28,†} M. Piccolo,²⁸ M. Rama,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹
M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹
G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹
U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. A. Nash,³²
M. B. Nikolich,³² W. Panduro Vazquez,³² X. Chai,³³ M. J. Charles,³³ U. Mallik,³³ N. T. Meyer,³³ V. Ziegler,³³
J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴
A. V. Gritsan,³⁵ M. Fritsch,³⁶ G. Schott,³⁶ N. Arnaud,³⁷ M. Davier,³⁷ G. Grosdidier,³⁷ A. Höcker,³⁷ F. Le
Diberder,³⁷ V. Lepeltier,³⁷ A. M. Lutz,³⁷ A. Oyanguren,³⁷ S. Pruvot,³⁷ S. Rodier,³⁷ P. Roudeau,³⁷ M. H. Schune,³⁷
A. Stocchi,³⁷ W. F. Wang,³⁷ G. Wormser,³⁷ C. H. Cheng,³⁸ D. J. Lange,³⁸ D. M. Wright,³⁸ C. A. Chavez,³⁹
I. J. Forster,³⁹ J. R. Fry,³⁹ E. Gabathuler,³⁹ R. Gamet,³⁹ K. A. George,³⁹ D. E. Hutchcroft,³⁹ D. J. Payne,³⁹
K. C. Schofield,³⁹ C. Touramanis,³⁹ A. J. Bevan,⁴⁰ F. Di Lodovico,⁴⁰ W. Menges,⁴⁰ R. Sacco,⁴⁰ G. Cowan,⁴¹
H. U. Flaecher,⁴¹ D. A. Hopkins,⁴¹ P. S. Jackson,⁴¹ T. R. McMahon,⁴¹ S. Ricciardi,⁴¹ F. Salvatore,⁴¹ A. C. Wren,⁴¹
D. N. Brown,⁴² C. L. Davis,⁴² J. Allison,⁴³ N. R. Barlow,⁴³ R. J. Barlow,⁴³ Y. M. Chia,⁴³ C. L. Edgar,⁴³
G. D. Lafferty,⁴³ M. T. Naisbit,⁴³ J. C. Williams,⁴³ J. I. Yi,⁴³ C. Chen,⁴⁴ W. D. Hulsbergen,⁴⁴ A. Jawahery,⁴⁴
C. K. Lae,⁴⁴ D. A. Roberts,⁴⁴ G. Simi,⁴⁴ G. Blaylock,⁴⁵ C. Dallapiccola,⁴⁵ S. S. Hertzbach,⁴⁵ X. Li,⁴⁵
T. B. Moore,⁴⁵ S. Saremi,⁴⁵ H. Staengle,⁴⁵ R. Cowan,⁴⁶ G. Sciolla,⁴⁶ S. J. Sekula,⁴⁶ M. Spitznagel,⁴⁶ F. Taylor,⁴⁶

R. K. Yamamoto,⁴⁶ H. Kim,⁴⁷ P. M. Patel,⁴⁷ S. H. Robertson,⁴⁷ A. Lazzaro,⁴⁸ V. Lombardo,⁴⁸ F. Palombo,⁴⁸ J. M. Bauer,⁴⁹ L. Cremaldi,⁴⁹ V. Eschenburg,⁴⁹ R. Godang,⁴⁹ R. Kroeger,⁴⁹ D. A. Sanders,⁴⁹ D. J. Summers,⁴⁹ H. W. Zhao,⁴⁹ S. Brunet,⁵⁰ D. Côté,⁵⁰ P. Taras,⁵⁰ F. B. Viaud,⁵⁰ H. Nicholson,⁵¹ N. Cavallo,^{52, ‡} G. De Nardo,⁵² F. Fabozzi,^{52, ‡} C. Gatto,⁵² L. Lista,⁵² D. Monorchio,⁵² P. Paolucci,⁵² D. Piccolo,⁵² C. Sciacca,⁵² M. Baak,⁵³ G. Raven,⁵³ H. L. Snoek,⁵³ C. P. Jessop,⁵⁴ J. M. LoSecco,⁵⁴ T. Allmendinger,⁵⁵ G. Benelli,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵ P. D. Jackson,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ A. M. Rahimi,⁵⁵ R. Ter-Antonyan,⁵⁵ Q. K. Wong,⁵⁵ N. L. Blount,⁵⁶ J. Brau,⁵⁶ R. Frey,⁵⁶ O. Igonkina,⁵⁶ M. Lu,⁵⁶ C. T. Potter,⁵⁶ R. Rahmat,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶ J. Strube,⁵⁶ E. Torrence,⁵⁶ F. Galeazzi,⁵⁷ A. Gaz,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ A. Pompili,⁵⁷ M. Posocco,⁵⁷ M. Rotondo,⁵⁷ F. Simonetto,⁵⁷ R. Stroili,⁵⁷ C. Voci,⁵⁷ M. Benayoun,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ L. Del Buono,⁵⁸ Ch. de la Vaissière,⁵⁸ O. Hamon,⁵⁸ B. L. Hartfiel,⁵⁸ M. J. J. John,⁵⁸ J. Malclès,⁵⁸ J. Ocariz,⁵⁸ L. Roos,⁵⁸ G. Therin,⁵⁸ P. K. Behera,⁵⁹ L. Gladney,⁵⁹ J. Panetta,⁵⁹ M. Biasini,⁶⁰ R. Covarelli,⁶⁰ C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹ F. Bucci,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ R. Cenci,⁶¹ F. Forti,⁶¹ M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹ M. A. Mazur,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ G. Rizzo,⁶¹ J. J. Walsh,⁶¹ M. Haire,⁶² D. Judd,⁶² D. E. Wagoner,⁶² J. Biesiada,⁶³ N. Danielson,⁶³ P. Elmer,⁶³ Y. P. Lau,⁶³ C. Lu,⁶³ J. Olsen,⁶³ A. J. S. Smith,⁶³ A. V. Telnov,⁶³ F. Bellini,⁶⁴ G. Cavoto,⁶⁴ A. D'Orazio,⁶⁴ D. del Re,⁶⁴ E. Di Marco,⁶⁴ R. Faccini,⁶⁴ F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ L. Li Gioi,⁶⁴ M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ G. Piredda,⁶⁴ F. Polci,⁶⁴ F. Safai Tehrani,⁶⁴ C. Voena,⁶⁴ M. Ebert,⁶⁵ H. Schröder,⁶⁵ R. Waldi,⁶⁵ T. Adye,⁶⁶ N. De Groot,⁶⁶ B. Franek,⁶⁶ E. O. Olaiya,⁶⁶ F. F. Wilson,⁶⁶ S. Emery,⁶⁷ A. Gaidot,⁶⁷ S. F. Ganzhur,⁶⁷ G. Hamel de Monchenault,⁶⁷ W. Kozanecki,⁶⁷ M. Legendre,⁶⁷ G. Vasseur,⁶⁷ Ch. Yèche,⁶⁷ M. Zito,⁶⁷ X. R. Chen,⁶⁸ H. Liu,⁶⁸ W. Park,⁶⁸ M. V. Purohit,⁶⁸ J. R. Wilson,⁶⁸ M. T. Allen,⁶⁹ D. Aston,⁶⁹ R. Bartoldus,⁶⁹ P. Bechtle,⁶⁹ N. Berger,⁶⁹ R. Claus,⁶⁹ J. P. Coleman,⁶⁹ M. R. Convery,⁶⁹ M. Cristinziani,⁶⁹ J. C. Dingfelder,⁶⁹ J. Dorfan,⁶⁹ G. P. Dubois-Felsmann,⁶⁹ D. Dujmic,⁶⁹ W. Dunwoodie,⁶⁹ R. C. Field,⁶⁹ T. Glanzman,⁶⁹ S. J. Gowdy,⁶⁹ M. T. Graham,⁶⁹ V. Halyo,⁶⁹ C. Hast,⁶⁹ T. Hryn'ova,⁶⁹ W. R. Innes,⁶⁹ M. H. Kelsey,⁶⁹ P. Kim,⁶⁹ D. W. G. S. Leith,⁶⁹ S. Li,⁶⁹ S. Luitz,⁶⁹ V. Luth,⁶⁹ H. L. Lynch,⁶⁹ D. B. MacFarlane,⁶⁹ H. Marsiske,⁶⁹ R. Messner,⁶⁹ D. R. Muller,⁶⁹ C. P. O'Grady,⁶⁹ V. E. Ozcan,⁶⁹ A. Perazzo,⁶⁹ M. Perl,⁶⁹ T. Pulliam,⁶⁹ B. N. Ratcliff,⁶⁹ A. Roodman,⁶⁹ A. A. Salnikov,⁶⁹ R. H. Schindler,⁶⁹ J. Schwiening,⁶⁹ A. Snyder,⁶⁹ J. Stelzer,⁶⁹ D. Su,⁶⁹ M. K. Sullivan,⁶⁹ K. Suzuki,⁶⁹ S. K. Swain,⁶⁹ J. M. Thompson,⁶⁹ J. Va'vra,⁶⁹ N. van Bakel,⁶⁹ M. Weaver,⁶⁹ A. J. R. Weinstein,⁶⁹ W. J. Wisniewski,⁶⁹ M. Wittgen,⁶⁹ D. H. Wright,⁶⁹ A. K. Yarritu,⁶⁹ K. Yi,⁶⁹ C. C. Young,⁶⁹ P. R. Burchat,⁷⁰ A. J. Edwards,⁷⁰ S. A. Majewski,⁷⁰ B. A. Petersen,⁷⁰ C. Roat,⁷⁰ L. Wilden,⁷⁰ S. Ahmed,⁷¹ M. S. Alam,⁷¹ R. Bula,⁷¹ J. A. Ernst,⁷¹ V. Jain,⁷¹ B. Pan,⁷¹ M. A. Saeed,⁷¹ F. R. Wappler,⁷¹ S. B. Zain,⁷¹ W. Bugg,⁷² M. Krishnamurthy,⁷² S. M. Spanier,⁷² R. Eckmann,⁷³ J. L. Ritchie,⁷³ A. Satpathy,⁷³ C. J. Schilling,⁷³ R. F. Schwitters,⁷³ J. M. Izen,⁷⁴ X. C. Lou,⁷⁴ S. Ye,⁷⁴ F. Bianchi,⁷⁵ F. Gallo,⁷⁵ D. Gamba,⁷⁵ M. Bomben,⁷⁶ L. Bosio,⁷⁶ C. Cartaro,⁷⁶ F. Cossutti,⁷⁶ G. Della Ricca,⁷⁶ S. Dittongo,⁷⁶ L. Lancieri,⁷⁶ L. Vitale,⁷⁶ V. Azzolini,⁷⁷ F. Martinez-Vidal,⁷⁷ Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ M. Pappagallo,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰ K. T. Flood,⁸⁰ J. J. Hollar,⁸⁰ P. E. Kutter,⁸⁰ B. Mellado,⁸⁰ A. Mihalys,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Universitat de Barcelona, Facultat de Física Dept. ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

- ¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA
- ¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA
- ¹⁹California Institute of Technology, Pasadena, California 91125, USA
- ²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
- ²¹University of Colorado, Boulder, Colorado 80309, USA
- ²²Colorado State University, Fort Collins, Colorado 80523, USA
- ²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
- ²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
- ²⁵Ecole Polytechnique, Laboratoire Leprince-Ringuet, F-91128 Palaiseau, France
- ²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- ²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ³⁰Harvard University, Cambridge, Massachusetts 02138, USA
- ³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³²Imperial College London, London, SW7 2AZ, United Kingdom
- ³³University of Iowa, Iowa City, Iowa 52242, USA
- ³⁴Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁵Johns Hopkins University, Baltimore, Maryland 21218, USA
- ³⁶Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
- ³⁷Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France
- ³⁸Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁹University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ⁴⁰Queen Mary, University of London, E1 4NS, United Kingdom
- ⁴¹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ⁴²University of Louisville, Louisville, Kentucky 40292, USA
- ⁴³University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴⁴University of Maryland, College Park, Maryland 20742, USA
- ⁴⁵University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴⁶Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁷McGill University, Montréal, Québec, Canada H3A 2T8
- ⁴⁸Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁹University of Mississippi, University, Mississippi 38677, USA
- ⁵⁰Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
- ⁵¹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁵²Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵³NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵⁴University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵⁵Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁶University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁷Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁸Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
- ⁵⁹University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁶⁰Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ⁶¹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶²Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶³Princeton University, Princeton, New Jersey 08544, USA
- ⁶⁴Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁵Universität Rostock, D-18051 Rostock, Germany
- ⁶⁶Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁷DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁸University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁹Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁷⁰Stanford University, Stanford, California 94305-4060, USA
- ⁷¹State University of New York, Albany, New York 12222, USA
- ⁷²University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷³University of Texas at Austin, Austin, Texas 78712, USA
- ⁷⁴University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷⁵Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁶Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁷IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
- ⁷⁸University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁹Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

⁸⁰University of Wisconsin, Madison, Wisconsin 53706, USA

⁸¹Yale University, New Haven, Connecticut 06511, USA

(Dated: February 6, 2008)

We report the first observation of e^+e^- annihilations into states of positive C -parity, namely $\rho^0\rho^0$ and $\phi\rho^0$. The two states are observed in the $\pi^+\pi^-\pi^+\pi^-$ and $K^+K^-\pi^+\pi^-$ final states, respectively, in a data sample of 225 fb^{-1} collected by the BABAR experiment at the PEP-II e^+e^- storage rings at energies near $\sqrt{s} = 10.58\text{ GeV}$. The distributions of $\cos\theta^*$, where θ^* is the center-of-mass polar angle of the ϕ meson or the forward ρ^0 meson, suggest production by two-virtual-photon annihilation. We measure cross sections within the range $|\cos\theta^*| < 0.8$ of $\sigma(e^+e^- \rightarrow \rho^0\rho^0) = 20.7 \pm 0.7(\text{stat}) \pm 2.7(\text{syst})\text{ fb}$ and $\sigma(e^+e^- \rightarrow \phi\rho^0) = 5.7 \pm 0.5(\text{stat}) \pm 0.8(\text{syst})\text{ fb}$.

PACS numbers: 12.20.-m, 12.38.Aw, 12.40.Vv

The process $e^+e^- \rightarrow \text{hadrons}$ at center-of-mass (CM) energy \sqrt{s} far below the Z^0 mass is dominated by annihilation via a single virtual photon with charge-conjugation parity $C = -1$. The high luminosity of the B factories provides an opportunity to explore rare, low multiplicity final states with $C = +1$ such as those produced in the two-virtual-photon annihilation (TVPA) process depicted in Fig. 1. The TVPA process has been ignored in the interpretation of the total hadronic cross section in e^+e^- annihilations as input to calculations [1] of the muon $g-2$ and the running QED coupling α . We report the first observation of the exclusive reactions $e^+e^- \rightarrow \rho^0\rho^0$ and $e^+e^- \rightarrow \phi\rho^0$, in which the final states are even under charge conjugation, and therefore cannot be produced via single-photon annihilation.

This analysis uses a 205 fb^{-1} data sample of e^+e^- collisions collected on the $\Upsilon(4S)$ resonance and 20 fb^{-1} collected 40 MeV below with the BABAR detector at the SLAC PEP-II asymmetric-energy B factory. The BABAR detector is described in detail elsewhere [2]. Charged-particle momenta and energy loss are measured in the tracking system which consists of a silicon vertex tracker (SVT) and a drift chamber (DCH). Electrons and photons are detected in a CsI(Tl) calorimeter (EMC). An internally reflecting ring-imaging Cherenkov detector (DIRC) provides charged particle identification (PID). An instrumented magnetic flux return (IFR) provides identification of muons. Kaon and pion candidates are identified using likelihoods of particle hypotheses calculated from the specific ionization in the DCH and SVT, and the Cherenkov angle measured in the DIRC. Electrons are identified by the ratio of the energy deposited in the EMC to the momentum and by the shower shape; muons are identified by the depth of penetration into the IFR.

Events with four well-reconstructed charged tracks and a total charge of zero are selected. Charged tracks are required to have at least 12 DCH hits and a polar angle in the range $0.41 < \theta < 2.54$ radians. The momenta of kaon and pion candidates are required to be greater than 800 and 600 MeV/c, respectively. Among the four selected tracks, two oppositely charged tracks must be identified as pions, and the other pair must be identified




FIG. 1: Two-virtual-photon annihilation diagram.

as two pions or two kaons. Events in which one or more pion candidates are identified as an electron or muon are rejected (lepton veto). We fit the four tracks to a common vertex, and require the χ^2 probability to exceed 0.1%. We accept events with reconstructed invariant mass within $170\text{ MeV}/c^2$ of the nominal CM energy (Fig. 2).

In the process $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$, there are two possible pairings of π^+ mesons with π^- mesons. However, only one combination appears in the kinematic region of interest ($m_{\pi^+\pi^-} < 2\text{ GeV}/c^2$) for both pairs. We label the pion pair with CM momentum vector pointing into the hemisphere defined by the e^- beam direction $\pi^+\pi^-_f$ and the other as $\pi^+\pi^-_b$. Figure 3(a) shows the scatter plot of the invariant masses of $\pi^+\pi^-_f$ and $\pi^+\pi^-_b$ from $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ events, and Fig. 3(b) the plot of invariant masses of K^+K^- and $\pi^+\pi^-$ pairs from $e^+e^- \rightarrow K^+K^-\pi^+\pi^-$ events. We observe correlations of masses in Fig. 3(a) indicating the production of $\rho^0\rho^0$ final states, and in Fig. 3(b) indicating the production of $\phi\rho^0$ final states.

To extract the number of $e^+e^- \rightarrow \rho^0\rho^0$ and $\phi\rho^0$ signal events, we perform a binned maximum-likelihood fit for nine rectangular regions (tiles) in the two-dimensional mass distributions, as shown in Fig. 3. The signal box is the central tile (tile 5), defined by the mass ranges $0.5 < m_{\pi^+\pi^-} < 1.1\text{ GeV}/c^2$ and $1.008 < m_{K^+K^-} < 1.035\text{ GeV}/c^2$. For $e^+e^- \rightarrow K^+K^-\pi^+\pi^-$, the expected number of events, n_i , for each tile i can be expressed as:




FIG. 2: Distributions of the invariant mass ($\Upsilon(4S)$ data) for the a) $\pi^+\pi^-\pi^+\pi^-$ and b) $K^+K^-\pi^+\pi^-$ final states. The accepted signal regions are indicated by the dashed lines.

$$n_i = f_i^S S + f_i^\phi N_\phi + f_i^{\rho^0} N_{\rho^0} + f_i^B B, \quad (1)$$

where S is the number of $\phi\rho^0$ signal events, N_ϕ is the number of ϕX background events, N_{ρ^0} is the number of $\rho^0 X$ background events, and B is the number of residual background events, in all nine tiles. The parameter f_i^T is the fraction of events of type T that contributes to tile i . The signal fractions f_i^S are modeled by Monte Carlo (MC) simulation [4], and f_i^ϕ and $f_i^{\rho^0}$ are obtained from the ϕX and $\rho^0 X$ background shapes, which are estimated by fitting the projections of $m_{K^+K^-}$ and $m_{\pi^+\pi^-}$ as described later. The residual background fractions f_i^B are modeled by a linear function that can be expressed as

$$f_i^B = \frac{\Delta x_i \Delta y_i [1 + s_{\rho^0}(x_i - x_5) + s_\phi(y_i - y_5)]}{\sum_{j=1}^9 \Delta x_j \Delta y_j}, \quad (2)$$

where Δx_i and Δy_i are the kinematically accessible dimensions of tile i , x_i and y_i are at the center of tile i , and s_{ρ^0} and s_ϕ are slopes obtained from the fits. A similar expression is used for the $\pi^+\pi^-\pi^+\pi^-$ case, where ϕ and ρ^0 are replaced with ρ_f^0 and ρ_b^0 .

The background fractions are obtained by mass projection fits which are confined to the central horizontal or central vertical ϕ or ρ^0 resonance band. The effect of neglecting the resonance width outside the central band, checked by smearing the background fractions in the central band into the adjacent tiles using the resonance widths obtained from MC, is found to be negligible. The mass projections in the central bands for $\pi^+\pi^-$ recoiling against a selected ρ^0 or ϕ and for K^+K^- recoiling against a ρ^0 are shown in Fig. 4. For the $\rho^0\pi^+\pi^-$ case we fit the $\pi^+\pi^-$ mass projection to the sum of a ρ^0 component, an $f_2(1270)$ component, and a $\mu^+\mu^-$ background component. The ρ^0 is represented by the product of a P-wave relativistic Breit-Wigner with its width set to the Particle Data Group (PDG) [3] value, a phase space term, and a factor $1/m_{\pi\pi}^2$ due to production via a virtual




FIG. 3: Scatter plots of the invariant masses of the two oppositely charged pairs in the a) $\pi^+\pi^-\pi^+\pi^-$ and b) $K^+K^-\pi^+\pi^-$ final states. The dashed lines indicate $K^+K^-/\pi^+\pi^-$ thresholds. The solid lines show the nine tiles used in the fit.

photon. The $f_2(1270)$ is represented by a D-wave relativistic Breit-Wigner with its mean and width set to the PDG values. The $\mu^+\mu^-$ background shape is obtained from a sample of the related channel $e^+e^- \rightarrow \rho^0\mu^+\mu^-$ isolated by requiring two oppositely charged tracks identified as muons. For the $\phi\pi^+\pi^-$ case, we use the same background parameterization in terms of f_2 and $\mu^+\mu^-$, but refit for their normalizations. For the $\rho^0 K^+K^-$ case, we fit the K^+K^- mass projection to the sum of a Breit-Wigner with mean and width fixed to their PDG values for the ϕ signal, and a threshold function $(q^3)/(1+q^3 R)$, where q is kaon momentum in the ϕ rest-frame and R is a shape parameter, for background. Assuming the masses of the two pairs to be uncorrelated and excluding the ρ^0 and ϕ signal contributions, the fitted functions are integrated to obtain the tile fractions $f_i^{\rho^0}$, f_i^ϕ , and $f_i^{\rho^0}$.

The extracted $\rho^0\rho^0$ and $\phi\rho^0$ yields in the signal box are 1243 ± 43 and 147 ± 13 events, to be compared with total of $1508 \pi^+\pi^-\pi^+\pi^-$ ($\sim 18\%$ background) and $163 K^+K^-\pi^+\pi^-$ ($\sim 10\%$ background) events in the signal box, respectively.

To investigate the possibility of $\rho^0\rho^0$ and $\phi\rho^0$ production in $\Upsilon(4S)$ decay, we examine the data recorded at and below the $\Upsilon(4S)$ resonance separately. The yields below the $\Upsilon(4S)$ resonance are 104 ± 14 for $\rho^0\rho^0$ and 14 ± 4 for $\phi\rho^0$, consistent with the expected values of 112 ± 4 and 13 ± 1 obtained by scaling the on-peak yields of 1138 ± 42 and 135 ± 13 by the relative integrated luminosities.

To investigate the production mechanism, we examine the production angle θ^* , defined as the angle between the ρ_f^0 (ϕ) direction and the e^- beam direction in the CM frame. To measure the angular distributions, we subdivide the data into bins of θ^* , and repeat the above fit, with linear background slopes $s_{\rho_f^0}$ and $s_{\rho_b^0}$ (s_{ρ^0} and s_ϕ) fixed to the values from the overall fit. The $|\cos\theta^*|$ distributions after MC efficiency correction are shown in Fig. 5. The measurements are restricted to the fiducial region $|\cos\theta^*| < 0.8$, as the efficiency drops rapidly be-




FIG. 4: Mass distribution for a) $\pi^+\pi^-$ pairs in $\rho^0\pi^+\pi^-$ events, b) $\pi^+\pi^-$ pairs in $\phi\pi^+\pi^-$ events, and c) K^+K^- pairs in $\rho^0K^+K^-$ events. The solid curves are the total fit. For the $\pi^+\pi^-$ cases, the dotted curve is the $\mu^+\mu^-$ component, while the sum of $f_2(1270)$ and $\mu^+\mu^-$ contributions are shown as dashed. For the K^+K^- case, the dashed curve represents the threshold function.

yond 0.8. These forward peaking $\cos\theta^*$ distributions are consistent with the TVPA expectation which we find can be approximated by:

$$\frac{d\sigma}{d\cos\theta^*} \propto \frac{1 + \cos^2\theta^*}{1 - \cos^2\theta^*} \quad (3)$$

in the fiducial region. The TVPA hypothesis gives a χ^2/dof (degrees of freedom) of 11.8/7 ($\rho^0\rho^0$) and 3.5/3 ($\phi\rho^0$). The fits disfavor $1 + \cos^2\theta^*$, giving a χ^2/dof of 112/7 for $\rho^0\rho^0$ and 6.3/3 for $\phi\rho^0$.

Other observables are the ϕ (ρ^0) decay helicity angles θ_H , defined as the angle, measured in the ϕ (ρ^0) rest frame, between the positively charged kaon or pion and the flight direction of the ϕ or ρ^0 in the CM frame. The efficiency-corrected distribution of $\cos\theta_H$, obtained using the procedure outline above for θ^* , is shown for the ρ^0 and ϕ candidates in Fig. 6. The solid lines in Fig. 6 are normalized $\sin^2\theta_H$ distributions which give χ^2/dof of 19.3/9 (ρ^0 from $\rho^0\rho^0$), 16.4/9 (ϕ from $\phi\rho^0$), and 3.1/9 (ρ^0 from $\phi\rho^0$). The $\sin^2\theta_H$ distributions indicate that ϕ and ρ^0 are transversely polarized as expected for TVPA. The dihedral angles, the angles between the decay planes of the two vector mesons measured in the CM frame, are consistent with a flat distribution with χ^2/dof of 7.0/9 ($\rho^0\rho^0$) and 10.9/9 ($\phi\rho^0$).

The combined hardware and software trigger efficiencies for signal events in the fiducial region are 99.9% for




FIG. 5: Production angle distributions, after correction for efficiency, for a) $\rho^0\rho^0$ and b) $\phi\rho^0$. The solid and dashed lines are the normalized $\frac{1+\cos^2\theta^*}{1-\cos^2\theta^*}$ and $1 + \cos^2\theta^*$ distributions, respectively.

$\rho^0\rho^0$ and 91.3% for $\phi\rho^0$. The lower efficiency for $\phi\rho^0$ is due to an event shape cut in the software trigger. For the determination of signal cross sections, the MC $\cos\theta^*$ and $\cos\theta_H$ distributions for ϕ and ρ^0 are re-weighted to reproduce the expectation from TVPA. The signal efficiencies in the fiducial region of $|\cos\theta^*| < 0.8$ for $\rho^0\rho^0$ and $\phi\rho^0$ are estimated to be 26.7% and 23.2%, respectively, including corrections to MC simulations of PID, tracking, hardware and software trigger efficiencies. Initial state photon radiation is included in the MC simulation.

Systematic uncertainties due to PID and tracking efficiency are estimated based on measurements from control data samples. The related systematic uncertainties on lepton vetoes are estimated by the difference from not applying the e and μ vetoes on pions. The systematic uncertainty from background subtraction is estimated by varying assumptions about background shapes. We investigated possible feed-down background from related modes with an extra π^0 using various extrapolations from the four-particle mass sidebands. We assume that the final states are fully transversely polarized. The systematic uncertainties are summarized in Table I.

Taking the branching fraction of $\phi \rightarrow K^+K^-$ as 49.1% and $\rho^0 \rightarrow \pi^+\pi^-$ as 100% [3], and signal mass regions of $0.5 < m_{\rho^0} < 1.1 \text{ GeV}/c^2$ and $1.008 < m_\phi < 1.035 \text{ GeV}/c^2$, we obtain the following results for the TVPA cross sections within $|\cos\theta^*| < 0.8$ near $\sqrt{s} = 10.58 \text{ GeV}$:

$$\begin{aligned} \sigma_{\text{fid}}(e^+e^- \rightarrow \rho^0\rho^0) &= 20.7 \pm 0.7(\text{stat}) \pm 2.7(\text{syst}) \text{ fb} \\ \sigma_{\text{fid}}(e^+e^- \rightarrow \phi\rho^0) &= 5.7 \pm 0.5(\text{stat}) \pm 0.8(\text{syst}) \text{ fb}. \end{aligned}$$

The measured cross sections are in good agreement with the calculation from a vector-dominance two-photon exchange model [7].

In summary, we have observed exclusive production of $C = +1$ final states in e^+e^- interactions. The measured C parity configuration, the signal yields in data samples on the $\Upsilon(4S)$ resonance and below, and the pro-




FIG. 6: Decay helicity angle distributions for a) ρ^0 from $\rho^0\rho^0$, b) ϕ from $\phi\rho^0$, c) ρ^0 from $\phi\rho^0$. The solid lines are the normalized $\sin^2\theta_H$ distributions.

TABLE I: Systematic uncertainties on the cross sections for $e^+e^- \rightarrow \rho^0\rho^0/\phi\rho^0$.

	$\rho^0\rho^0$	$\phi\rho^0$
Particle Identification	9.6%	10.4%
Background subtraction	7.0%	7.0%
Tracking efficiency	5.0%	5.0%
$\rho^0\rho^0\pi^0, \phi\rho^0\pi^0$ background	1.6%	2.7%
Luminosity	1.2%	1.2%
Total	13.0%	14.0%

duction angle distributions support the conclusion that the production mechanism is two-virtual-photon annihilation. The Standard Model predictions of the anomalous magnetic moment of the muon and the QED coupling rely on the measurements of low-energy e^+e^- hadronic cross sections, which are assumed to be entirely due to single-photon exchange. We have estimated the effect due to the TVPA processes we have measured, and find it to be small compared with the current precision [1].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. We wish to thank S. Brodsky, M. Peskin and H. Quinn for helpful discussions. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONA-CyT (Mexico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France

† Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡ Also with Università della Basilicata, Potenza, Italy

- [1] M. Davier, S. Eidelman, A. Hoecker, Z. Zhang, *Eur. Phys. J. C* **31**, 503 (2003).
- [2] *BABAR* Collaboration, B. Aubert *et al.*, *Nucl. Instrum. Methods, Sect. A* **479**, 1 (2002).
- [3] Particle Data Group, S. Eidelman *et al.*, *Phys. Lett. B* **592**, 1 (2004).
- [4] $e^+e^- \rightarrow \rho^0\rho^0/\phi\rho^0$ are generated uniformly over phase space. We use the AFKQED package to simulate the signal processes, including hard and multiple soft initial state radiation, using the methods in [5] and [6].
- [5] H. Czyż and J.H. Kühn, *Eur. Phys. J. C* **18**, 497 (2001).
- [6] M. Caffo, H. Czyż, and E. Remiddi, *Nuo. Cim.* **110A**, 515 (1997); *Phys. Lett. B* **327**, 369 (1994).
- [7] M. Davier, M. Peskin, and A. Snyder, hep-ph/0606155.